

**RPB465Ra01 100µg**

**Recombinant Connexin 40 (CX40)**

**Organism Species: Rattus norvegicus (Rat)**

***Instruction manual***

FOR IN VITRO USE AND RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

---

---

11th Edition (Revised in May, 2016)

## **[ PROPERTIES ]**

**Source:** Prokaryotic expression.

**Host:** *E. coli*

**Residues:** Ala50~Glu303

**Tags:** N-terminal His-Tag

**Tissue Specificity:** Lung.

**Subcellular Location:** Cell membrane; Multi-pass membrane protein. Cell junction, gap junction.

**Purity:** >98%

**Traits:** Freeze-dried powder

**Buffer formulation:** 20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% sarcosyl, 5%Trehalose and Proclin300.

**Original Concentration:** 200ug/mL

**Applications:** SDS-PAGE; WB; ELISA; IP; CoIP; Purification; Amine Reactive Labeling.

(May be suitable for use in other assays to be determined by the end user.)

**Predicted isoelectric point:** 6.9

**Predicted Molecular Mass:** 30.1kDa

**Accurate Molecular Mass:** 20&16kDa as determined by SDS-PAGE reducing conditions.

**Phenomenon explanation:**

The possible reasons that the actual band size differs from the predicted are as follows:

1. Splice variants: Alternative splicing may create different sized proteins from the same gene.
2. Relative charge: The composition of amino acids may affects the charge of the protein.
3. Post-translational modification: Phosphorylation, glycosylation, methylation etc.
4. Post-translation cleavage: Many proteins are synthesized as pro-proteins, and then cleaved to give the active form.
5. Polymerization of the target protein: Dimerization, multimerization etc.

## **[ USAGE ]**

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

## **[ STORAGE AND STABILITY ]**

**Storage:** Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

## **[ SEQUENCE ]**

A

```
DFRCDTIQPG CQNVCYDQAF PISHIRYWVL QIIFVSTPSL VYMGHAMHTV
RMQEKQKLRE AEKAKEAGGT GTYEYLAEKA ELSCWKEVNG KIVLQGTLLN
TYVCTILIRT AMEVAFMVGQ YLLYGIFLDT LHVCRRSPCP HPVNCYVSRP
TEKNVFIVFM MAVAGLSLFL SLAELYHLGW KKIRQRLAKS RQGDKHQLLG
PSTSLVQGLT PPPDFNQCLK NSPDEKFFSD FSNNMGSRKN PDPLATEEVP
NQE
```

