

RPB231Ra01 10µg

Recombinant Cytokeratin 18 (CK18)

Organism Species: Rattus norvegicus (Rat)

Instruction manual

FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)



[PROPERTIES]

Source: Prokaryotic expression

Host: E.coli

Residues: Glu72~Gly380

Tags: N-terminal His Tag

Subcellular Location: Nucleus, Cytoplasm

Purity: > 97%

Traits: Freeze-dried powder

Buffer formulation: 20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT,

0.01% SKL, 5% Trehalose and Proclin300.

Original Concentration: 200µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 5.2

Predicted Molecular Mass: 39.6kDa

Accurate Molecular Mass: 40kDa as determined by SDS-PAGE reducing conditions.

[USAGE]

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCE]

		EKETMQDLN	DRLASYLDKV	KNLETENRRL	
ESKIREYLEK	RGPQGVRDWG	HYFKTIEDLR	AQIFANSVDN	ARIVLQIDNA	
RLAADDFRVK	YETELAMRQS	VESDIHGLRK	VVDDTNITRL	QLETEIEALK	
EELLFMKKNH	EEEVQGLEAQ	IASSGLTVEV	DAPKSQDLSK	IMADIRAQYE	
QLAQKNREEL	DKYWSQQIEE	STTVVTTKSA	EIRDAETTLL	ELRRTLQTLE	
IDLDSMKNQN	INLENNLGEV	EARYRVQMEQ	LNGVLLHLES	ELAQTRAEGQ	
RQTQEYEALL	NIKVKLEAEI	ATYRRLLEDG			

[IDENTIFICATION]




Figure . Gene Sequencing (extract)




Figure. SDS-PAGE

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.