

RPG444Mu01 50µg

Recombinant Dipeptidyl Peptidase 9 (DPP9)

Organism Species: Mus musculus (Mouse)

Instruction manual

FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)



[PROPERTIES]

Source: Prokaryotic expression

Host: E.coli

Residues: Val505~Leu862

Tags: N-terminal His and GST Tag

Subcellular Location: Cytoplasm

Purity: > 90%

Traits: Freeze-dried powder

Buffer formulation: 100mMNaHCO₃, 500mMNaCl, pH8.3, containing 0.01% SKL, 5%

Trehalose.

Original Concentration: 200µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 6.3

Predicted Molecular Mass: 70.7kDa

Accurate Molecular Mass: 71kDa as determined by SDS-PAGE reducing conditions.

[USAGE]

Reconstitute in ddH₂O to a concentration of 0.1-1.0 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCE]

```
VNEQTK LVYFQGTKDT PLEHHLYVVS YESAGEIVRL TTLGFSHSCS
MSQSFDMFVS HYSSVSTPPC VHVYKLSGPD DDPLHKQPRF WASMMEAANC
PPDYVPPEIF HFHTRADVQL YGMIYKPHTL QPGRKHPTVL FVYGGPQVQL
VNNSFKGIKY LRLNTLASLG YAVVVIDGRG SCQRGLHFEG ALKNQMGQVE
IEDQVEGLQY VAEKYGFIDL SRVAIHGWSY GGFLSLMGLI HKPQVFKVAI
AGAPVTVWMA YDTGYTERYM DVPENNQQGY EAGSVALHVE KLPNEPNRLL
ILHGFLDENV HFFHTNFLVS QLIRAGKPYQ LQIYPNERHS IRCRESGEHY
EVTLLHFLQE HL
```

[IDENTIFICATION]




Figure . Gene Sequencing (extract)




Figure. SDS-PAGE

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.