

RPJ114Hu01 10µg

Recombinant Fc Fragment Of IgG Receptor Transporter Alpha (FcgRT)

Organism Species: Homo sapiens (Human)

Instruction manual

FOR RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

12th Edition (Revised in Aug, 2016)



[PROPERTIES]

Source: Prokaryotic expression

Host: E.coli

Residues: Ala24~Ala365

Tags: N-terminal His Tag

Subcellular Location: Membrane

Purity: > 90%

Traits: Freeze-dried powder

Buffer formulation: 20mM Tris, 150mM NaCl, pH8.0, containing 0.01% skl, 5%Trehalose.

Original Concentration: 50µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 5.9

Predicted Molecular Mass: 41.0kDa

Accurate Molecular Mass: 41kDa as determined by SDS-PAGE reducing conditions.

[USAGE]

Reconstitute in ddH₂O to a concentration of 0.1-0.5 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.


Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCE]



		AESHLSL	LYHLTAVSSP	APGTPAFWVS
GWLGPQQYLS	YNSLRGEAEP	CGAWVWENQV	SWYWEKETTD	LRIKEKLFLE
AFKALGGKGP	YTLQGLLGCE	LGPDNTSVPT	AKFALNGEEF	MNFDLKQGTW
GGDWPEALAI	SQRWQQQDKA	ANKELTFLLF	SCPHRLREHL	ERGRGNLEWK
EPPSMRLKAR	PSSPGFSVLT	CSAFSFYPPE	LQLRFLRNGL	AAGTGQGDFG
PNSDGSFHAS	SSLTVKSGDE	HHYCCIVQHA	GLAQPLRVEL	ESPAKSSVLV
VGIVIGVLLL	TAAAVGGALL	WRRMRSGLPA	PWISLRGDDT	GVLLPTPGEA
QDADLKDVNV	IPATA			

[IDENTIFICATION]



[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.