

RPD022Bo01 100ug

Recombinant Fibrinogen Like Protein 1 (FGL1)

Organism Species: Bos taurus; Bovine (Cattle)

Instruction manual

FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)



[PROPERTIES]

Source: Prokaryotic expression

Host: E.coli

Residues: Leu23~Val312 (Accession # Q3SZZ7)

Tags: N-terminal His Tag

Subcellular Location: Secreted

Purity: > 90%

Traits: Freeze-dried powder

Buffer formulation: PBS, pH7.4, containing 0.01% SKL, 5% Trehalose.

Original Concentration: 200µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 5.96

Predicted Molecular Mass: 40.4kDa

Accurate Molecular Mass: 40/18kDa as determined by SDS-PAGE reducing conditions.

[USAGE]

Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.


Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCE]




LENCLQEQ ARLRAQVYLL ETRVKQQQVK ISQLLHEKQV QLLDKGEENS VIDLGGKRQY ADCSEIFNDG YKQSGFYKIK PLQSPAEFSV YCDMSDGGGW TVIQRRSDGS ENFNRDWSDY ENGFGNFVQK NGEYWLGNRN LHLLTTQGDY TLKIDLADFE KNSRYAQYKN FKVGDEKNSY DLHIGEYSGT AGDSLTGNFH PEVQWWASHQ RMKFSTWDRDNDNYEGNCAKEDQSGWWFNRCHSANL NGFY HKGPYTAKTD NGIVWHTWHG WWYSLKSVVM KIRPNDFIPN IV

[IDENTIFICATION]



CTOS MAN CTROCTOCAGS MOST COSCICAGE CONSIGNATOR CTROCTOS MOST CAME ACCESS AND ACCESS AND

Figure. Gene Sequencing (Extract)



[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.