RPC482Ra01 100ug Recombinant Flavin Containing Monooxygenase 2, Non Functional (FMO2) **Organism Species: Rattus norvegicus (Rat)** Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

Coud-Clone Corp.

10th Edition (Revised in Jan, 2014)


[PROPERTIES]

Residues: Arg116~Val324 **Tags:** Two N-terminal Tags, His-tag and T7-tag Accession: Q6IRI9 Host: E. coli Subcellular Location: Microsome Membrane. Endoplasmic Reticulum Membrane. **Purity:** >95% Endotoxin Level: <1.0EU per 1µg (determined by the LAL method). Formulation: Supplied as lyophilized form in 20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% sarcosyl, 5% trehalose, and preservative. 15% SDS-PAGE Predicted isoelectric point: 8.6 Predicted Molecular Mass: 27.8kDa Applications: SDS-PAGE; WB; ELISA; IP.

(May be suitable for use in other assays to be determined by the end user.)

[USAGE]

Reconstitute in sterile ddH₂O.





[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

[<u>SEQUENCES</u>]

The sequence of the target protein is listed below.

RPDFA SSGQWDVYVQ SNGKEQRAVF DAVMVCSGHH IQPHLPLKSF PGIERFQGQY FHSRQYKHPV GYEGKRILVV GIGNSAADIA SELSKRAAQV FVSTRHGSWV LSRISEDGYP WDMVFHTRFS SMLRNVLPRT VVKWMMERQM NRWFNHENYG LVPQNKYLMK EPVLNDDLPS RLLYGAIKVK TRVKELTETA VVFEDGTVEE DVDV