

RPF195Hu01 1 Recombinant Fucosyltransferase 5 (FUT5) Organism Species: *Homo sapiens (Human) Instruction manual*

FOR RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

12th Edition (Revised in Aug, 2016)

Coud-Clone Corp.

[PROPERTIES]

Source: Prokaryotic expression

Host: E.coli

Residues: Arg35~Thr374

Tags: N-terminal His Tag

Subcellular Location: Secreted

Purity: > 90%

Traits: Freeze-dried powder

Buffer formulation: 20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT,

0.01% SKL, 5% Trehalose and Proclin300.

Original Concentration: 200µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point:

Predicted Molecular Mass: 40.4kDa

Accurate Molecular Mass: 43kDa as determined by SDS-PAGE reducing conditions.

[<u>USAGE</u>]

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.



[<u>SEQUENCE</u>]

			RVSRDD	ATGSPRPGLM
AVEPVTGAPN	GSRCQDSMAT	PAHPTLLILL	WTWPFNTPVA	LPRCSEMVPG
AADCNITADS	SVYPQADAVI	VHHWDIMYNP	SANLPPPTRP	QGQRWIWFSM
ESPSNCRHLE	ALDGYFNLTM	SYRSDSDIFT	PYGWLEPWSG	QPAHPPLNLS
AKTELVAWAV	SNWKPDSARV	RYYQSLQAHL	KVDVYGRSHK	PLPKGTMMET
LSRYKFYLAF	ENSLHPDYIT	EKLWRNALEA	WAVPVVLGPS	RSNYERFLPP
DAFIHVDDFQ	SPKDLARYLQ	ELDKDHARYL	SYFRWRETLR	PRSFSWALAF
CKACWKLQQE	SRYQTVRSIA	AWFT		

[IDENTIFICATION]




Figure. SDS-PAGE

[<u>IMPORTANT NOTE</u>]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.