

RPA308Bo01 100µg Recombinant Galectin 8 (GAL8) Organism Species: *Bos taurus; Bovine (Cattle) Instruction manual* 

FOR RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

12th Edition (Revised in Aug, 2016)

# Cloud-Clone Corp.

### [PROPERTIES]

Source: Prokaryotic expression

Host: E.coli

Residues: Met1~Trp357

Tags: N-terminal His Tag

Subcellular Location: Secreted

**Purity:** > 90%

Traits: Freeze-dried powder

Buffer formulation: 20mM Tris, 150mM NaCl, pH8.0, containing 0.01% SKL, 5% Trehalose.

Original Concentration: 200µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 9.0

Predicted Molecular Mass: 43.8kDa

Accurate Molecular Mass: 41kDa as determined by SDS-PAGE reducing conditions.

#### [<u>USAGE</u>]

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

#### [ STORAGE AND STABILITY ]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

#### [ <u>SEQUENCE</u> ]

# Cloud-Clone Corp.

MMSLNNLQNV IYNPVIPYVG TISEQLEPGT LIVLRGHVPS DSDRFQVDLQ CGSSVKPRAD VAFHFNPRFK RANCVVCNTL RNEKWGWEEI TYDMPFKKEK SFEIVIMVLK EKFQVAVNGR HTLLYAHRIS PERIDTLGIY GKVIIHSVGF SFSSDLGSTQ GSTLEPTGIS KENVQKSGES QLPSNRGDIS KIVPRTVYTK SKASPANHTL TCAKILPTNC LSKTLPFVAR LNSSMGPGRT IVIKGEVNTN AKGFTVDLLS GKSKDIALHL NPRLNVKAFV RNSFLQEAWG EEERNITCFP FSPGMYFEMI IYCDAREFKV AVNGVHSLEY KHRFKELSKV DTLEIDGDIH LLEVRSW

### [IDENTIFICATION]




Figure. SDS-PAGE

#### [IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.