

RPA309Ca01 100µg

**Recombinant Galectin 9 (GAL9)** 

Organism Species: Canis familiaris; Canine (Dog)

Instruction manual

FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

12th Edition (Revised in Aug, 2016)



### [PROPERTIES]

**Source:** Prokaryotic expression

Host: E.coli

Residues: Phe17~Thr355

Tags: N-terminal His Tag

Subcellular Location: n/a

**Purity:** > 95%

Traits: Freeze-dried powder

Buffer formulation: 20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT,

0.01% SKL, 5% Trehalose and Proclin300.

Original Concentration: 200µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 9.1

Predicted Molecular Mass: 51.7kDa

Accurate Molecular Mass: 53kDa as determined by SDS-PAGE reducing conditions.

#### [USAGE]

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

### [STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.



# [SEQUENCE]

	FSGK	IQGGLQDGLK	ITINGTILYC	NGTRFAVNFH
SGHSDSDIAF	HFNPRFEEGG	YVVCNTKQKG	SWGSEERKMQ	MPFQMGNPFE
LCFMVNSCDF	KVTVNGSHFT	QYSHRVPFHY	VDTLSITGAV	QLSYISFQNT
RVTITQPAFS	TMQFPQTACF	PPRPKGRKPK	SPGVWQSSSA	PITQTVIHTV
QTTPGQPFPN	PIIPPTAYPT	PTYPMPFFTS	<b>IPGGLYPSKS</b>	IIVSGTVLPG
AKRFHINLRS	GNDIAFHLNP	RFNENTVVRN	TQINNSWGSE	ERSLPRKMPF
VQGQSFSVWI	MCEGHCFKVA	VDGEHLFEYY	HRLKNLLAIN	NMEVAGDVQL
THVQT				

# [ IDENTIFICATION ]




Figure. SDS-PAGE

# [ IMPORTANT NOTE ]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.