

RPJ051Hu01 100ug

Recombinant Guanine Deaminase (GDA)

Organism Species: *Homo sapiens (Human)*

Instruction manual

FOR RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)

[PROPERTIES]

Source: Prokaryotic expression

Host: *E.coli*

Residues: Met1~Val454

Tags: N-terminal His Tag

Subcellular Location: Plasma

Purity: > 80%

Traits: Freeze-dried powder

Buffer formulation: PBS, pH7.4, containing 0.01% SKL, 5% Trehalose.

Original Concentration: 200µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 5.4

Predicted Molecular Mass: 54.7kDa

Accurate Molecular Mass: 55kDa as determined by SDS-PAGE reducing conditions.

[USAGE]

Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.


Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCE]

MCAAQMPPLA HIFRGTFVHS TWTCPMEVLR DHLLGVSDSG KIVFLEEASQ
QEKLAKWCF KPCEIRELSH HEFFMPGLVD THIHASQYSF AGSSIDLPLL
EWLTKYTFPA EHRFQNI DFA EEVYTRVRR TLKNGTTTAC YFATIHTDSS
LLLADITDKF GQRAFVGKVC MDLNDTFPEY KETTEESIKE TERFVSEMLQ
KNYSRVKPIV TPRFSLSCSE TLMGELGNIA KTRDLHIQSH ISENREVEA
VKNLYPSYKN YTSVYDKNNL LTNKTVMAHG CYLSAEELNV FHERGASIAH
CPNSNLSLSS GFLNVLEV LK HEVKIGLCTD VAGGYSYSML DAIRRAVMVS
NILLINKVNE KSLTLKEVFR LATLGGSQAL GLDGEIGNFE VGKEFDAILI
NPKASDSPID LFYGDFFGDI SEAVIQKFLY LGDDRNIEEV YVGGKQVVPF
SSSV

[IDENTIFICATION]



[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.