

RPB995Hu02 100µg Recombinant Hemojuvelin (HJV) Organism Species: *Homo sapiens (Human)* Instruction manual

FOR RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)

# Coud-Clone Corp.

### [PROPERTIES]

Source: Prokaryotic expression
Host: E.coli

Residues: Arg41~Ala386

Tags: N-terminal His Tag

Subcellular Location: Membrane

**Purity:** > 95%

Traits: Freeze-dried powder

Buffer formulation: PBS, pH7.4, containing 0.01% SKL, 5% Trehalose .

Original Concentration: 200µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 8.3

Predicted Molecular Mass: 40.3kDa

Accurate Molecular Mass: 43kDa as determined by SDS-PAGE reducing conditions.

#### [ <u>USAGE</u> ]

Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

#### [ STORAGE AND STABILITY ]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

#### [SEQUENCE]

## Cond-Clone Corp.

				RCNAEYVSST	LSLRGGGSSG
ALRGGGGGGR	GGGVGSGGLC	RALRSYALCT	RRTARTCRGD	LAFHSAVHGI	EDLMIQHNCS
RQGPTAPPPP	RGPALPGAGS	GLPAPDPCDY	EGRF SRLHGR	PPGFLHCASF	GDPHVRSFHH
HFHTCRVQGA	WPLLDNDFLF	VQATSSPMAL	GANATATRKL	TIIFKNMQEC	IDQKVYQAEV
DNLPVAFEDG	SINGGDRPGG	SSLSIQTANP	GNHVEIQAAY	IGTTIIIRQT	AGQLSFSIKV
AEDVAMAFSA	EQDLQLCVGG	CPPSQRLSRS	ERNRRGAITI	DTARRLCKEG	LPVEDAYFHS
CVFDVLISGD	PNFTVAAQAA	LEDARA			

## [ IDENTIFICATION ]





Figure. Gene Sequencing (Extract)



## [<u>IMPORTANT NOTE</u>]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.