

RPL617Hu01 10µg Recombinant Insulin Like Growth Factor 2 mRNA Binding Protein 3 (IGF2BP3) Organism Species: *Homo sapiens (Human)* Instruction manual

FOR RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

12th Edition (Revised in Aug, 2016)

Coud-Clone Corp.

[PROPERTIES]

Source: Prokaryotic expression

Host: E.coli

Residues: Asn2~lle343

Tags: N-terminal His Tag

Subcellular Location: Nucleus, Cytoplasm

Purity: > 95%

Traits: Freeze-dried powder

Buffer formulation: 20mM Tris, 500mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT,

0.01% SKL, 5% Trehalose and Proclin300.

Original Concentration: 200µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 8.5

Predicted Molecular Mass: 41.6kDa

Accurate Molecular Mass: 42kDa as determined by SDS-PAGE reducing conditions.

[<u>USAGE</u>]

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.


Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.



[<u>SEQUENCE</u>]

NKLYIGNLS	ENAAPSDLES	IFKDAKIPVS	GPFLVKTGYA	FVDCPDESWA
LKAIEALSGK	IELHGKPIEV	EHSVPKRQRI	RKLQIRNIPP	HLQWEVLDSL
LVQYGVVESC	EQVNTDSETA	VVNVTYSSKD	QARQALDKLN	GFQLENFTLK
VAYIPDEMAA	QQNPLQQPRG	RRGLGQRGSS	RQGSPGSVSK	QKPCDLPLRL
LVPTQFVGAI	IGKEGATIRN	ITKQTQSKID	VHRKENAGAA	EKSITILSTP
EGTSAACKSI	LEIMHKEAQD	IKFTEEIPLK	ILAHNNFVGR	LIGKEGRNLK
KIEQDTDTKI	TISPLQELTL	YNPERTITVK	GNVETCAKAE	EEI

[IDENTIFICATION]



[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.