

RPC093Hu01 1 Recombinant Integrin Alpha 7 (ITGa7) Organism Species: *Homo sapiens (Human) Instruction manual* 

FOR RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

12th Edition (Revised in Aug, 2016)

# Cond-Clone Corp.

## [PROPERTIES]

Source: Prokaryotic expression

Host: E.coli

Residues: Arg648~Ser952

Tags: N-terminal His Tag

Subcellular Location: Membrane, Secreted

**Purity:** > 90%

Traits: Freeze-dried powder

Buffer formulation: 20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT,

0.01% SKL, 5% Trehalose and Proclin300.

Original Concentration: 200µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 5

Predicted Molecular Mass: 36.9kDa

Accurate Molecular Mass: 37kDa as determined by SDS-PAGE reducing conditions.

#### [<u>USAGE</u>]

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

#### [ STORAGE AND STABILITY ]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.



## [SEQUENCE]

RQA

```
PGQGLPPVAP ILNAHQPSTQ RAEIHFLKQG CGEDKICQSN LQLVRARFCT
RVSDTEFQPL PMDVDGTTAL FALSGQPVIG LELMVTNLPS DPAQPQADGD
DAHEAQLLVM LPDSLHYSGV RALDPAEKPL CLSNENASHV ECELGNPMKR
GAQVTFYLIL STSGISIETT ELEVELLLAT ISEQELHPVS ARARVFIELP
LSIAGMAIPQ QLFFSGVVRG ERAMQSERDV GSKVKYEVTV SNQGQSLRTL
GSAFLNIMWP HEIANGKWLL YPMQVELEGG QGPGQKGLCS PRPNILHLDV
DS
```

### [IDENTIFICATION]




Figure. SDS-PAGE

#### [IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.