

RPB586Hu03 200µg

Recombinant Integrin Beta 2 (CD18)

Organism Species: Homo sapiens (Human)

Instruction manual

FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)



## [PROPERTIES]

**Source:** Prokaryotic expression

Host: E.coli

Residues: Gln23~Val367

Tags: N-terminal His Tag

**Subcellular Location:** Membrane

**Purity:** > 90%

Traits: Freeze-dried powder

**Buffer formulation:** PBS, pH7.4, containing 0.01% SKL, 5% Trehalose .

Original Concentration: 80µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 6.6

Predicted Molecular Mass: 42.0kDa

**Accurate Molecular Mass:** 44kDa as determined by SDS-PAGE reducing conditions.

#### [USAGE]

Reconstitute in ddH<sub>2</sub>O to a concentration of 0.1-0.5 mg/mL. Do not vortex.

### [STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

### [ SEQUENCE ]



		DECTVEVU	CCCDECTECC	DCCTLICOVI N
		QECINFRY	SSCRECIESG	PUCTWCQKLN
FTGPGDPDSI	RCDTRPQLLM	RGCAADDIMD	PTSLAETQED	HNGGQKQLSP
QKVTLYLRPG	QAAAFNVTFR	RAKGYPIDLY	YLMDLSYSML	DDLRNVKKLG
GDLLRALNEI	TESGRIGFGS	<b>FVDKTVLPFV</b>	NTHPDKLRNP	CPNKEKECQP
PFAFRHVLKL	TNNSNQFQTE	VGKQLISGNL	DAPEGGLDAM	MQVAACPEEI
<b>GWRNVTRLLV</b>	FATDDGFHFA	GDGKLGAILT	PNDGRCHLED	NLYKRSNEFD
YPSVGQLAHK	LAENNIQPIF	AVTSRMVKTY	EKLTEIIPKS	AVGELSEDSS
NVVQLIKNAY	NKLSSRV			

## [IDENTIFICATION]




Figure. SDS-PAGE

# [ IMPORTANT NOTE ]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.