

RPB598Mu01 1

**Recombinant Interferon Regulatory Factor 5 (IRF5)** 

Organism Species: Mus musculus (Mouse)

Instruction manual

FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

12th Edition (Revised in Aug, 2016)



## [ PROPERTIES ]

**Source:** Prokaryotic expression

Host: E.coli

Residues: His3~Glu250

Tags: N-terminal His Tag

Subcellular Location: Nucleus, Cytoplasm

**Purity:** > 95%

Traits: Freeze-dried powder

Buffer formulation: 100mMNaHCO<sub>3</sub>, 500mMNaCl, pH8.3, containing 1mM EDTA, 1mM DTT,

0.01% SKL, 5% Trehalose and Proclin300.

Original Concentration: 200µg/mL

**Applications:** Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 5.5

Predicted Molecular Mass: 31.1kDa

Accurate Molecular Mass: 34kDa as determined by SDS-PAGE reducing conditions.

#### [USAGE]

Reconstitute in 100mM NaHCO<sub>3</sub>, 500mM NaCl (pH8.3) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

### [STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.



# [SEQUENCE]

HSAPGIPP	PPRRVRLKPW	LVAQVNSCQY	PGLQWVNGEK	KLFYIPWRHA
TRHGPSQDGD	NTIFKAWAKE	TGKYTEGVDE	ADPAKWKANL	RCALNKSRDF
QLFYDGPRDM	PPQPYKIYEV	CSNGPAPTES	QPTDDYVLGE	EEEEEEELQ
RMLPGLSITE	PALPGPPNAP	YSLPKEDTKW	PPALQPPVGL	GPPVPDPNLL
APPSGNPAGE	ROLL PEVLEP	GPLASSOPPT	<b>FPLIPDLITS</b>	PHMI PLITDLE

## [ IDENTIFICATION ]




Figure . Gene Sequencing (extract)




Figure. SDS-PAGE

# [ IMPORTANT NOTE ]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.