

RPA386Hu02 10µg

Recombinant Lipoprotein lipase (LPL)

Organism Species: Homo sapiens (Human)

Instruction manual

FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

12th Edition (Revised in Aug, 2016)



[PROPERTIES]

Source: Prokaryotic expression

Host: E.coli

Residues: Ala28~Gly475

Tags: N-terminal His Tag

Subcellular Location: Membrane, Secreted

Purity: > 90%

Traits: Freeze-dried powder

Buffer formulation: PBS, pH7.4, containing 0.01% SKL, 5% Trehalose.

Original Concentration: 200µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 8.1

Predicted Molecular Mass: 54.1kDa

Accurate Molecular Mass: 54kDa as determined by SDS-PAGE reducing conditions.

[USAGE]

Reconstitute in ddH₂O to a concentration of 0.1-0.5 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCE]

		ADQ	RRDFIDIESK	FALRTPEDTA
EDTCHLIPGV	AESVATCHFN	HSSKTFMVIH	GWTVTGMYES	WVPKLVAALY
KREPDSNVIV	VDWLSRAQEH	YPVSAGYTKL	VGQDVARFIN	WMEEEFNYPL
DNVHLLGYSL	GAHAAGIAGS	LTNKKVNRIT	GLDPAGPNFE	YAEAPSRLSP
DDADFVDVLH	TFTRGSPGRS	IGIQKPVGHV	DIYPNGGTFQ	PGCNIGEAIR
VIAERGLGDV	DQLVKCSHER	SIHLFIDSLL	NEENPSKAYR	CSSKEAFEKG
LCLSCRKNRC	NNLGYEINKV	RAKRSSKMYL	KTRSQMPYKV	FHYQVKIHFS
GTESETHTNQ	AFEISLYGTV	AESENIPFTL	PEVSTNKTYS	FLIYTEVDIG
ELLMLKLKWK	SDSYFSWSDW	WSSPGFAIQK	IRVKAGETQK	KVIFCSREKV
SHLQKGKAPA	VFVKCHDKSL	NKKSG		

[IDENTIFICATION]




Figure. SDS-PAGE

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.