

RPB856Hu02 50µg Recombinant Major Histocompatibility Complex Class I G (MHCG) Organism Species: *Homo sapiens (Human) Instruction manual*

FOR RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

12th Edition (Revised in Aug, 2016)

Cond-Clone Corp.

[PROPERTIES]

Source: Prokaryotic expression

Host: E.coli

Residues: Gly25~lle308

Tags: N-terminal His Tag

Subcellular Location: Membrane, Secreted

Purity: > 90%

Traits: Freeze-dried powder

Buffer formulation: 100mMNaHCO₃, 500mMNaCl, pH8.3, containing 1mM EDTA, 1mM DTT,

0.01% SKL, 5% Trehalose and Proclin300.

Original Concentration: 200µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 5.0

Predicted Molecular Mass: 36.4kDa

Accurate Molecular Mass: 38kDa as determined by SDS-PAGE reducing conditions.

[<u>USAGE</u>]

Reconstitute in ddH_2O to a concentration of 0.1-1.0 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[<u>SEQUENCE</u>]

Cond-Clone Corp.

GSHSMR YFSAAVSRPG RGEPRFIAMG YVDDTQFVRF DSDSACPRME PRAPWVEQEG PEYWEEETRN TKAHAQTDRM NLQTLRGYYN QSEASSHTLQ WMIGCDLGSD GRLLRGYEQY AYDGKDYLAL NEDLRSWTAA DTAAQISKRK CEAANVAEQR RAYLEGTCVE WLHRYLENGK EMLQRADPPK THVTHHPVFD YEATLRCWAL GFYPAEIILT WQRDGEDQTQ DVELVETRPA GDGTFQKWAA VVVPSGEEQR YTCHVQHEGL PEPLMLRWKQ SSLPTIPI

[IDENTIFICATION]




Figure . Gene Sequencing (extract)




Figure. SDS-PAGE

[<u>IMPORTANT NOTE</u>]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.