

RPG192Ra01 50µg Recombinant N-Acetyltransferase 2 (NAT2) Organism Species: *Rattus norvegicus (Rat) Instruction manual* 

FOR RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)

# Coud-Clone Corp.

# [PROPERTIES]

Source: Prokaryotic expression

Host: E.coli

Residues: Asp20~Val280

Tags: N-terminal His and GST Tag

Subcellular Location: Cytoplasm

**Purity:** > 95%

Traits: Freeze-dried powder

Buffer formulation: 100mMNaHCO<sub>3</sub>, 500mMNaCl, pH8.3, containing 1mM DTT, 0.01% SKL,

5% Trehalose .

Original Concentration: 200µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 5.1

Predicted Molecular Mass: 60.3kDa

Accurate Molecular Mass: 60kDa as determined by SDS-PAGE reducing conditions.

#### [<u>USAGE</u>]

Reconstitute in  $ddH_2O$  to a concentration of 0.1-1.0 mg/mL. Do not vortex.

### [ STORAGE AND STABILITY ]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

## [<u>SEQUENCE</u>]



D LEELTEILQH QIRAIPFENL NIHCGESMEL NLEVIFDQVV RKKRGGWCLQ VNHLLYWALT KMGFEATMLG GYVFNTPANK YSSGMIHLLV QVTLSGKDYI VDAGFGRSYQ MWEPLELTSG KDQPQVPAIF RLTEENGTWY LDQIRREQYV PNQEFVNSDL LEKNKYRKIY SFTLEPRTIE DFESINTYLQ TSPASLFTSK SFCSLQTLEG VHCLVGSTLT YRRFSYKDNI DLVEFKSLTE EEIEDVLKTI FGVSLERKLV

#### [IDENTIFICATION]




Figure. SDS-PAGE

### [IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.