

RPA127Hu01 10µg

Recombinant Protease Activated Receptor 4 (PAR4)

Organism Species: Homo sapiens (Human)

Instruction manual

FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)



[PROPERTIES]

Source: Prokaryotic expression

Host: E.coli

Residues: Gly48~Gln385

Tags: N-terminal His and GST Tag

Subcellular Location: Membrane

Purity: > 97%

Traits: Freeze-dried powder

Buffer formulation: 20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT,

0.01% SKL, 5% Trehalose .

Original Concentration: 200µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 8.6

Predicted Molecular Mass: 66.3kDa

Accurate Molecular Mass: 66kDa as determined by SDS-PAGE reducing conditions.

[USAGE]

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.



[SEQUENCE]

				GYP
GQVCANDSDT	LELPDSSRAL	LLGWVPTRLV	PALYGLVLVV	GLPANGLALW
VLATQAPRLP	STMLLMNLAA	ADLLLALALP	PRIAYHLRGQ	RWPFGEAACR
LATAALYGHM	YGSVLLLAAV	SLDRYLALVH	PLRARALRGR	RLALGLCMAA
WLMAAALALP	LTLQRQTFRL	ARSDRVLCHD	ALPLDAQASH	WQPAFTCLAL
LGCFLPLLAM	LLCYGATLHT	LAASGRRYGH	ALRLTAVVLA	SAVAFFVPSN
LLLLLHYSDP	SPSAWGNLYG	AYVPSLALST	LNSCVDPFIY	YYVSAEFRDK
VRAGLFORSP	GDTVASKASA	EGGSRGMGTH	SSLL0	

[IDENTIFICATION]




Figure . Gene Sequencing (extract)




Figure. SDS-PAGE

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.