

RPQ311Hu01 100µg

Recombinant Proteasome 26S Subunit, Non ATPase 6 (PSMD6)

Organism Species: Homo sapiens (Human)

Instruction manual

FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)



[PROPERTIES]

Source: Prokaryotic expression

Host: E.coli

Residues: Met1~Met389

Tags: N-terminal His and GST Tag

Subcellular Location: Nucleus, Cytoplasm, Extracellular matrix

Purity: > 97%

Traits: Freeze-dried powder

Buffer formulation: PBS, pH7.4, containing 0.01% SKL, 5% Trehalose .

Original Concentration: 200µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 5.5

Predicted Molecular Mass: 75.5kDa

Accurate Molecular Mass: 76kDa as determined by SDS-PAGE reducing conditions.

[USAGE]

Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.


Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCE]



```
MPLENLEEEG LPKNPDLRIA QLRFLLSLPE HRGDAAVRDE LMAAVRDNNM
APYYEALCKS LDWQIDVDLL NKMKKANEDE LKRLDEELED AEKNLGESEI
RDAMMAKAEY LCRIGDKEGA LTAFRKTYDK TVALGHRLDI VFYLLRIGLF
YMDNDLITRN TEKAKSLIEE GGDWDRRNRL KVYQGLYCVA IRDFKQAAEL
FLDTVSTFTS YELMDYKTFV TYTVYVSMIA LERPDLREKV IKGAEILEVL
HSLPAVRQYL FSLYECRYSV FFQSLAVVEQ EMKKDWLFAP HYRYYVREMR
IHAYSQLLES YRSLTLGYMA EAFGVGVEFI DQELSRFIAA GRLHCKIDKV
NEIVETNRPD SKNWQYQETI KKGDLLLNRV QKLSRVINM
```

[IDENTIFICATION]



[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.