

RPB532Mu01 10µg

Recombinant Retinoic Acid Inducible Gene 1 Protein (RIG1)

Organism Species: Mus musculus (Mouse)

Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

11th Edition (Revised in May, 2016)

[PROPERTIES]

Source: Prokaryotic expression.

Host: E. coli

Residues: Met1~Gln211
Tags: N-terminal His-Tag

Tissue Specificity: Liver, Spleen, Lung, Heart.

Subcellular Location: Cell projection, ruffle membrane. Cytoplasm, cytoskeleton.

Cell junction, tight junction.

Purity: >95%

Endotoxin Level: <1.0EU per 1µg (determined by the LAL method).

Traits: Freeze-dried powder

Buffer formulation: 20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA,

1mM DTT, 0.01% sarcosyl, 5%Trehalose and Proclin300.

Original Concentration: 200ug/mL

Applications: SDS-PAGE; WB; ELISA; IP; CoIP; ReporterAssays; Purification;

Amine Reactive Labeling.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 5.2

Predicted Molecular Mass: 28.4kDa

Accurate Molecular Mass: 28kDa as determined by SDS-PAGE reducing conditions.



[USAGE]

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCE]

MTAEQRQNLQ AFRDYIKKIL DPTYILSYMS SWLEDEEVQY IQAEKNNKGP MEAASLFLQY LLKLQSEGWF QAFLDALYHA GYCGLCEAIE SWDFQKIEKL EEHRLLLRRL EPEFKATVDP NDILSELSEC LINQECEEIR QIRDTKGRMA GAEKMAECLI RSDKENWPKV LQLALEKDNS KFSELWIVDK GFKRAESKAD EDDGAEASSI Q

[IDENTIFICATION]




Figure 1. SDS-PAGE