

RPE969Hu01 100ug Recombinant Ring Finger Protein 1 (RNF1) Organism Species: *Homo sapiens (Human)* Instruction manual

FOR RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)

# Cond-Clone Corp.

## [PROPERTIES]

Source: Prokaryotic expression Host: *E.coli* Residues: Met1~Lys406 Tags: N-terminal His Tag Subcellular Location: Nucleus Purity: > 90% Traits: Freeze-dried powder Buffer formulation: pH7.4, containing 0.01% SKL, 5% Trehalose Original Concentration: 200µg/mL Applications: Positive Control; Immunogen; SDS-PAGE; WB. (May be suitable for use in other assays to be determined by the end user.) Predicted isoelectric point: 5.4 Predicted Molecular Mass: 46.1kDa

Accurate Molecular Mass: 43kDa as determined by SDS-PAGE reducing conditions.

## [ <u>USAGE</u> ]

Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex

#### [ STORAGE AND STABILITY ]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.


**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

#### [SEQUENCE]

## Cond-Clone Corp.

MTTPANAQNA SKTWELSLYE LHRTPQEAIM DGTEIAVSPR SLHSELMCPI CLDMLKNTM TTKECLHRFC SDCIVTALRS GNKECPTCRK KLVSKRSLRP D PNFDALIS KIYPSREEYE AHQDRVLIRL SRLHNQQALS SSIEEGLRMQ AM HRAQRVR RPIPGSDQTT TMSGGEGEPG EGEGDGEDVS SDSAPDSAPG PAP KRPRGG GAGGSSVGTG GGGTGGVGGG AGSEDSGDRG GTLGGGTLGP PSPP GAPSP PEPGGEIELV FRPHPLLVEK GEYCQTRYVK TTGNATVDHL SKYLA LRIA LERRQQQEAG EPGGPGGGAS DTGGPDGCGG EGGGAGGGDG PEEPAL PSL EGVSEKQYTI YIAPGGGAFT TLNGSLTLEL VNEKFWKVSR PLELCYA PT KDPK

### [IDENTIFICATION]



#### [IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.