

RPB843Hu01 200µg Recombinant Signaling Lymphocytic Activation Molecule Family, Member 2 (SLAMF2) Organism Species: *Homo sapiens (Human) Instruction manual*

FOR RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

12th Edition (Revised in Aug, 2016)

Cloud-Clone Corp.

[PROPERTIES]

Source: Prokaryotic expression

Host: E.coli

Residues: Gln27~Thr243

Tags: N-terminal His Tag

Subcellular Location: Membrane

Purity: > 97%

Traits: Freeze-dried powder

Buffer formulation: 100mMNaHCO₃, 500mMNaCl, pH8.3, containing 0.01% SKL, 5%

Trehalose.

Original Concentration: 200µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 8.7

Predicted Molecular Mass: 28.6kDa

Accurate Molecular Mass: 29kDa as determined by SDS-PAGE reducing conditions.

[<u>USAGE</u>]

Reconstitute in 100mM NaHCO3, 500mM NaCl (pH8.3) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Cond-Clone Corp.

[SEQUENCE]

QGHL VHMTVVSGSN VTLNISESLP ENYKQLTWFY TFDQKIVEWD SRKSKYFESK FKGRVRLDPQ SGALYISKVQ KEDNSTYIMR VLKKTGNEQE WKIKLQVLDP VPKPVIKIEK IEDMDDNCYL KLSCVIPGES VNYTWYGDKR PFPKELQNSV LETTLMPHNY SRCYTCQVSN SVSSKNGTVC LSPPCTLARS FGVEWIASWL VVTVPTILGL LLT

[IDENTIFICATION]





Figure. Gene Sequencing (Extract)



[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.