

RPB155Hu01 50µg

Recombinant Signaling Lymphocytic Activation Molecule Family, Member 5 (SLAMF5)

Organism Species: *Homo sapiens (Human)*

Instruction manual

FOR RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

12th Edition (Revised in Aug, 2016)

[**PROPERTIES**]

Source: Prokaryotic expression

Host: *E.coli*

Residues: Ile43~Leu248

Tags: N-terminal His Tag

Subcellular Location: Membrane, Chromosome

Purity: > 97%

Traits: Freeze-dried powder

Buffer formulation: 100mMNaHCO₃, 500mMNaCl, pH8.3, containing 0.01% SKL, 5% Trehalose.

Original Concentration: 500µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 6.8

Predicted Molecular Mass: 27.0kDa

Accurate Molecular Mass: 29kDa as determined by SDS-PAGE reducing conditions.

[**USAGE**]

Reconstitute in ddH₂O to a concentration of 0.1-1.0 mg/mL. Do not vortex.

[**STORAGE AND STABILITY**]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.


Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[**SEQUENCE**]

IIAWTSKTSV AYVTPGDSET APVVTVTHRN YYERIHALGP NYNLVISDLR IQEPRQVK
MEDAGDYKAD INTQADPYTT TKRYNLQIYR RLGKPKITQS LMASVNSTCN
VTLTCSVEKE EKNVTYNWSP LGEEGNVLQI FQTPEDQELT YTCTAQNPVS
NNSDSISARQ LCADIAMGFR THHTGLLSVL AMFFLLVLIL SSVFLFRL

[IDENTIFICATION]



[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.