

RPE698Hu01 200µg Recombinant Sphingosine Kinase 1 (SPHK1) Organism Species: *Homo sapiens (Human) Instruction manual*

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

12th Edition (Revised in Aug, 2016)

Coud-Clone Corp.

[PROPERTIES]

Source: Prokaryotic expression

Host: E.coli

Residues: Leu148~Leu398

Tags: N-terminal His Tag

Subcellular Location: Nucleus, Cytoplasm

Purity: > 90%

Traits: Freeze-dried powder

Buffer formulation: 20mM Tris, 150mM NaCl, pH8.0, containing 0.01% SKL, 5% Trehalose

and Proclin300.

Original Concentration: 200µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 6.9

Predicted Molecular Mass: 29.2kDa

Accurate Molecular Mass: 32kDa as determined by SDS-PAGE reducing conditions.

[<u>USAGE</u>]

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.


Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Cond-Clone Corp.

[<u>SEQUENCE</u>]

LT NCTLLLCRRL LSPMNLLSLH TASGLRLFSV LSLAWGFIAD VDLESEKYRR LGEMRFTLGT FLRLAALRTY RGRLAYLPVG RVGSKTPASP VVVQQGPVDA HLVPLEEPVP SHWTVVPDED FVLVLALLHS HLGSEMFAAP MGRCAAGVMH LFYVRAGVSR AMLLRLFLAM EKGRHMEYEC PYLVYVPVVA FRLEPKDGKG VFAVDGELMV SEAVQGQVHP NYFWMVSGCV EPPPSWKPQQ MPPPEEPL

[IDENTIFICATION]



[IMPORTANT NOTE]

The kit is designed for in vitro and research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.