

RPG483Hu01 10µg

Recombinant Tubulointerstitial Nephritis Antigen Like Protein 1 (TINAGL1)

Organism Species: *Homo sapiens (Human)*

Instruction manual

FOR RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

12th Edition (Revised in Aug, 2016)

[PROPERTIES]

Source: Prokaryotic expression

Host: *E.coli*

Residues: Val81~His467

Tags: N-terminal His Tag

Subcellular Location: Secreted

Purity: > 97%

Traits: Freeze-dried powder

Buffer formulation: 20mM Tris, 150mM NaCl, pH8.0, containing 0.01% SKL, 5% Trehalose.

Original Concentration: 200µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 6.6

Predicted Molecular Mass: 47.3kDa

Accurate Molecular Mass: 47kDa as determined by SDS-PAGE reducing conditions.

[USAGE]

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCE]

```

                                VSDCCPDFWD FCLGVPPFP
PIQGCMHGGR IYPVLGTYWD NCNRCTCQEN RQWQCDQEP LVDPMIKAI
NQGNYGWQAG NHSAFWGML DEGIRYRLGT IRPSSVMNM HEIYTVLNP
EVLPTAFEAS EKWPNIHEP LDQNCAGSW AFSTAAVSD RVSISLGHM
TPVLSQNL SCDTHQQGC RGGRLDGAW FLRRRGVSD HCYPFSGRER
DEAGPAPCM MHSRAMGRK RQATAHCPNS YVNNNDIYQV TPVYRLGSND
KEIMKELMEN GPVQALMEVH EDFFLYKGGI YSHTPVSLGR PERYRRHGTH
SVKITGWGEE TLPDGRTLKY WTAANSWGPA WGERGHFRIV RGVNECDIES
FVLGVWGRVG MEDMGHH

```

[IDENTIFICATION]




Figure. SDS-PAGE

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.