

RPC829Hu01 100µg Recombinant Tuftelin (TUFT) Organism Species: *Homo sapiens (Human)* Instruction manual

FOR RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)

# Cond-Clone Corp.

### [PROPERTIES]

Source: Prokaryotic expression

Host: E.coli

Residues: Met1~Thr390

Tags: N-terminal His and GST Tag

Subcellular Location: Secreted

**Purity:** > 90%

**Traits:** Freeze-dried powder

Buffer formulation: PBS, pH7.4, containing 0.01% SKL, 5% Trehalose.

Original Concentration: 200µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 5.7

Predicted Molecular Mass: 74.3kDa

Accurate Molecular Mass: 74kDa as determined by SDS-PAGE reducing conditions.

### [<u>USAGE</u>]

Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

#### [ STORAGE AND STABILITY ]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

### [SEQUENCE]

## Cloud-Clone Corp.

MNGTRNWCTL	VDVHPEDQAA	GSVDILRLTL	QGELTGDELE	HIAQKAGRKT
YAMVSSHSAG	HSLASELVES	HDGHEEIIKV	YLKGRSGDKM	IHEKNINQLK
SEVQYIQEAR	NCLQKLREDI	SSKLDRNLGD	SLHRQEIQVV	LEKPNGFSQS
PTALYSSPPE	VDTCINEDVE	SLRKTVQDLL	AKLQEAKRQH	QSDCVAFEVT
LSRYQREAEQ	SNVALQREED	RVEQKEAEVG	ELQRRLLGME	TEHQALLAKV
REGEVALEEL	RSNNADCQAE	REKAATLEKE	VAGLREKIHH	LDDMLKSQQR
KVRQMIEQLQ	NSKAVIQSKD	ATIQELKEKI	AYLEAENLEM	HDRMEHLIEK
QISHGNFSTQ	ARAKTENPGS	IRISKPPSPK	PMPVIRVVET	

### [IDENTIFICATION]

### [<u>IMPORTANT NOTE</u>]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.