RPF805Mu01 10µg

Recombinant Tumor Necrosis Factor Alpha Induced Protein 3 Interacting Protein 2

(TNIP2)

Organism Species: *Mus musculus (Mouse)*

Instruction manual

FOR RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

12th Edition (Revised in Aug, 2016)

[PROPERTIES]

Source: Prokaryotic expression

Host: *E.coli*

Residues: Ala170~Phe413

Tags: N-terminal His Tag

Subcellular Location: Cytoplasm

Purity: > 95%

Traits: Freeze-dried powder

Buffer formulation: 20mM Tris, 150mM NaCl, pH8.0, containing 0.01% SKL, 5% Trehalose.

Original Concentration: 200µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 6.6

Predicted Molecular Mass: 31.9kDa

Accurate Molecular Mass: 33kDa as determined by SDS-PAGE reducing conditions.

[USAGE]

Reconstitute in ddH$_2$O to a concentration of 0-0.1mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8⁰C for one month.

Aliquot and store at -80⁰C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCE]

```
            A KCLDERQCAQ GDAGEKSPAE LEQTSSDASG
QSVIKKLQEE NRLLKQKVTH VEDLNAKWQR YDASRDEYVK GLHAQLKRRQ
VPLEPELMKK EISRLNRQLE EKISDCAEAN QELTAMRMSR DTALERVQML
EQQILAYKDD FKSERADRER AHSRIQELEE KIMSLMYQVS QRQDSREPGP
CRIHTGNKTA KYLEMDALEH VTPGGWRPES RSQQMEPSAE GGHVCTAQRG
QGDLQCPHCL RCF
```

[IDENTIFICATION]

Figure . Gene Sequencing (extract)

15% SDS-PAGE

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.