

RPA139Ca01 50µg

Recombinant Tumor Necrosis Factor Related Apoptosis Inducing Ligand (TRAIL)

Organism Species: Canis familiaris; Canine (Dog)

Instruction manual

FOR RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)



[PROPERTIES]

Source: Prokaryotic expression

Host: E.coli

Residues: Thr36~Gly281

Tags: N-terminal His Tag

Subcellular Location: Secreted

Purity: > 90%

Traits: Freeze-dried powder

Buffer formulation: PBS, pH7.4, containing 0.01% SKL, 1mM DTT, 5% Trehalose and

Proclin300.

Original Concentration: 1200µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 8.6

Predicted Molecular Mass: 32.6kDa

Accurate Molecular Mass: 33kDa as determined by SDS-PAGE reducing conditions.

[USAGE]

Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCE]



| | | | TRELK | QMQDKYSQSG |
|------------|------------|------------|------------|------------|
| IACFLKEDDI | PWDPSDEESM | NNPCWQVKWQ | LRQFVRKMIL | KTYEETIPTA |
| PEKQLNIPYV | VSDRGSQRVA | AHITGTSRRS | MFPIPSSKND | KALGHKINSW |
| DSTRKGHSFL | NNLHLRNGEL | VIHQRGFYYI | YSQTYFRFQE | PEEIPTGQNR |
| KRNKQMVQYI | YKHTSYPDPI | LLMKSARNSC | WSKDSEYGLY | SIYQGGIFEL |
| KENDRIFVSV | SNEQLIDMDQ | EASFFGAFLI | G | |

[IDENTIFICATION]




Figure. SDS-PAGE

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.