

**RPB367Hu01 1mg** 

Recombinant Vascular Endothelial Growth Factor Receptor 2 (VEGFR2)

Organism Species: Homo sapiens (Human)

Instruction manual

FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)



## [PROPERTIES]

**Source:** Prokaryotic expression

Host: E.coli

Residues: Asn46~Thr320

Tags: N-terminal His Tag

Subcellular Location: Membrane, Nucleus, Cytoplasm, Endosome

**Purity:** > 80%

Traits: Freeze-dried powder

**Buffer formulation:** PBS, pH7.4, containing 0.01% Sarcosyl, 5% Trehalose.

Original Concentration: 300µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 7.6

Predicted Molecular Mass: 34.6kDa

**Accurate Molecular Mass:** 35kDa as determined by SDS-PAGE reducing conditions.

#### [USAGE]

Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

### [STORAGE AND STABILITY]

**Storage:** Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

### [ SEQUENCE ]

				NTTLQ
ITCRGQRDLD	WLWPNNQSGS	<b>EQRVEVTECS</b>	DGLFCKTLTI	PKVIGNDTGA
YKCFYRETDL	ASVIYVYVQD	YRSPFIASVS	DQHGVVYITE	NKNKTVVIPC
LGSISNLNVS	LCARYPEKRF	VPDGNRISWD	SKKGFTIPSY	MISYAGMVFC
EAKINDESYQ	SIMYIVVVVG	YRIYDVVLSP	SHGIELSVGE	KLVLNCTART
ELNVGIDFNW	EYPSSKHQHK	KLVNRDLKTQ	SGSEMKKFLS	TLTIDGVTRS
<b>DQGLYTCAAS</b>	SGLMTKKNST			

# [ IDENTIFICATION ]




Figure . Gene Sequencing (extract)




Figure. SDS-PAGE

## [ IMPORTANT NOTE ]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.