

RPP358Hu01 50µg

Recombinant Wingless Type MMTV Integration Site Family, Member 10A (WNT10A)

Organism Species: Homo sapiens (Human)

Instruction manual

FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

12th Edition (Revised in Aug, 2016)



# [PROPERTIES]

**Source:** Prokaryotic expression

Host: E.coli

Residues: Ser108~Pro343

Tags: N-terminal His and GST Tag

Subcellular Location: Secreted

**Purity:** > 97%

Traits: Freeze-dried powder

Buffer formulation: PBS, pH7.4, containing 0.01% SKL, 1mM DTT, 5% Trehalose and

Proclin300.

Original Concentration: 200µg/mL

**Applications:** Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 10.5

Predicted Molecular Mass: 56.0kDa

Accurate Molecular Mass: 56kDa as determined by SDS-PAGE reducing conditions.

### [USAGE]

Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

#### [ STORAGE AND STABILITY ]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.


Aliquot and store at -80°C for 12 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

### [ SEQUENCE ]


```
SSL ETRNKIPYES PIFSRGFRES AFAYAIAAAG VVHAVSNACA
LGKLKACGCD ASRRGDEEAF RRKLHRLQLD ALQRGKGLSH GVPEHPALPT
ASPGLQDSWE WGGCSPDMGF GERFSKDFLD SREPHRDIHA RMRLHNNRVG
RQAVMENMRR KCKCHGTSGS CQLKTCWQVT PEFRTVGALL RSRFHRATLI
RPHNRNGGQL EPGPAGAPSP APGAPGPRRR ASPADLVYFE KSP
```

## [ IDENTIFICATION ]



### ANAMATION TACAGA GARDON AT THE ACCORDENT THE CALCAGA GARDON TACAGA G

Figure . Gene Sequencing (extract)



## [ IMPORTANT NOTE ]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.