

RPA948Ra01 50µg

Recombinant Thymidine Phosphorylase (TP)

Organism Species: Rattus norvegicus (Rat)

kDa 70 Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

10th Edition (Revised in Jan, 2014)

[PROPERTIES]

Residues: Ile33~Ala290

Tags: Two N-terminal Tags, His-tag and GST-tag

Accession: Q5FVR2

Host: E. coli

Purity: >90%

Endotoxin Level: <1.0EU per 1µg

(determined by the LAL method).

Formulation: Supplied as lyophilized form in 20mM Tris,

150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT,

0.01% sarcosyl, 5% trehalose, and preservative.

Predicted isoelectric point: 5.8

Predicted Molecular Mass: 59.2kDa

Applications: SDS-PAGE; WB; ELISA; IP.

(May be suitable for use in other assays to be determined by the end user.)

[USAGE]

Reconstitute in sterile ddH₂O.



[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCES]

The sequence of the target protein is listed below.

IRLKRNGG HLSEADIRNF VHALMDGRAQ DTQIGAMLMA IRLQGMDLEE TSVLTQALAE SGQQLEWPKA WHQQLVDKHS TGGVGDKVSL VLAPALAACG CKVPMISGRS LGHTGGTLDK LESIPGFSVT QSPEQMLQIL EEVGCCIVGQ SEKLVPADGI LYAARDVTAT VDSVPLITAS ILSKKAVEGL STLVVDVKFG GAAVFPDQEK ARELAKMLVR VGMGLGLQVA AALTAMDNPL GRNVGHTLEV EEALLCLDGA

[REFERENCES]

- 1. Kubiak R., et al. (1999) Z. Naturforsch., C, J. Biosci. 54:1096-1102.
- 2. Zhao B., et al. (2000) Jpn. J. Cancer Res. 91:331-336.
- 3. Yanagi Y., et al. (2003) Invest. Ophthalmol. Vis. Sci. 44:751-754.
- 4. Miszczak-Zaborska E., et al. (2004) Gynecol. Oncol. 94:86-92.