



A92705Hu01 Polyclonal Antibody to Oxytocin Receptor (OXTR) Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

5th Edition (Revised in January, 2013)

[PRODUCT INFORMATION]

Immunogen: OXTR **Clonality:** Polyclonal

Host: Rabbit


Species Reactivity: Human

Ig type: Rabbit IgG

Purification: Antibodies are purified by target protein affinity chromatography.

Applications: WB, ICC, IHC-P, IHC-F

Form: Liquid Size: 100µg



Sample: Recombinant human OXTR

[IMMUNOGEN INFORMATION]

Immunogen: Recombinant human OXTR (Arg272~Ser377) expressed in E.coli.

Molecular Weight: 39.2kDa

USCN accession No.: P92705Hu01

Sequence: The target protein is fused with two N-terminal Tags, His-tag and

GST-tags-Tag and its sequence is listed below.

MRNKKFELGL EFPNLPYYID GDVKLTQSMA IIRYIADKHN MLGGCPKERA EISMLEGAVL DIRYGVSRIA YSKDFETLKV DFLSKLPEML KMFEDRLCHK TYLNGDHVTH PDFMLYDALD VVLYMDPMCL DAFPKLVCFK KRIEAIPQID KYLKSSKYIA WPLQGWQATF GGGDHPPKSD GSTSGSGHHH HHHSAGLVPR GSTAIGMKET AAAKFERQHM DSPDLGTLEV LFQ





GPLGSEF- RTVKMTFII VLAFIVCWTP FFFVQMWSVW DANAPKEASA FIIVMLLASL NSCCNPWIYM LFTGHLFHEL VQRFLCCSAS YLKGRRLGET SASKKSNSSS FVLSHRS

[ANTIBODY SPEFICITY]

Anti OXTR is a rabbit polyclonal antibody raised against human OXTR. This antibody has been selected for its ability to recognize human OXTR in immunohistochemical staining and western blotting, non cross-reactive with other members of the family.

[APPLICATIONS]

Western blotting: 1:100-400

Immunocytochemistry in formalin fixed cells: 1:100-500

Immunohistochemistry in formalin fixed frozen section: 1:100-500

Immunohistochemistry in paraffin section: 1:50-200

Optimal working dilutions must be determined by end user.

[CONTENTS]

Form & Buffer: Supplied as solution form in PBS, pH 7.4, containing 0.02%Na₃N, 50% glycerol.

[QUALITY CONTROL]

Content: The quality control contains recombinant human OXTR (Arg272~Ser377) disposed in loading buffer.

Usage: 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate. 5uL per well when used in enhanced chemilumescent (ECL).

Note: The quality control is specifically manufactured as the positive control. Not used for other purposes.

Loading buffer: 100mM Tris(PH 8.8), 2%SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, Na₃N 0.02%.

[STORAGE]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.