

EPA757Hu61 5mg
Eukaryotic Epidermal Growth Factor Receptor (EGFR)
Organism Species: *Homo sapiens (Human)*
Instruction manual

FOR RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)

[PROPERTIES]

Source: Eukaryotic expression

Host: 293F cell

Residues: Leu25~Ser645

Tags: N-terminal His Tag

Subcellular Location: Membrane

Purity: > 95%

Traits: Freeze-dried powder

Buffer formulation: PBS, pH7.4, containing 5% Trehalose.

Original Concentration: 400µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 6.5

Predicted Molecular Mass: 70.2kDa

Accurate Molecular Mass: 90kDa as determined by SDS-PAGE reducing conditions.

Phenomenon explanation:

The possible reasons that the actual band size differs from the predicted are as follows:

1. Splice variants: Alternative splicing may create different sized proteins from the same gene.
2. Relative charge: The composition of amino acids may affect the charge of the protein.
3. Post-translational modification: Phosphorylation, glycosylation, methylation etc.
4. Post-translation cleavage: Many proteins are synthesized as pro-proteins, and then cleaved to give the active form.
5. Polymerization of the target protein: Dimerization, multimerization etc.

[USAGE]

Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[**SEQUENCE**]

```

                LEEKKV CQGTSNKLTQ LGTFEDHFLS
LQRMFNCEV VLG NLEITYV QRNYDLSFLK TIQEVAGYVL IALNTVERIP
LENLQIIRGN MYYENSYALA VLSNYDANKT GLKELPMRNL QEILHGAVRF
SNPALCNEVE SIQWRDIVSS DFLSNMSMDF QNHLGSCQKC DPSCPNGSCW
GAGEENCQKL TKIICAQQCS GRCRGKSPSD CCHNQCAAGC TGPRESDECLV
CRKFRDEATC KDTCPPLMLY NPTTYQMDVN PEGKYSFGAT CVKKCPRNYV
VTDHGSCVRA CGADSYEMEE DGVRKCKKCE GPCRKVCNGI GIGEFKDSLS
INATNIKHFK NCTSISGLH ILPVAFRGDS FTHTPPLDPQ ELDILKTVKE
ITGFLLIQAW PENRTDLHAF ENLEIIRGRT KQHGQFSLAV VSLNITSLGL
RSLKEISDGD VIISGNKNLC YANTINWKKL FGTSGQKTKI ISNRGENSCK
ATGQVCHALC SPEGCWGPEP RDCVSCRNVS RGRECVDKCN LLEGEPREFV
ENSECIQCHP ECLPQAMNIT CTGRGPDNCI QCAHYIDGPH CVKTCAPAGVM
GENNTLVWKY ADAGHVCHLC HPNCTYGCTG PGLEGCP TNG PKIPS
    
```

[**IDENTIFICATION**]

CTGCGGAAAGGAAAGCTTTCCAGGCGCAGGTACCAAGCTGACCGCTTGGGCGCTTTGGAGATCATTTTCTGACCTGCGAGGAGGTTGATGATCTTTCTCTTAAAGGACATCCAGGAGTGGCTGGTTATGCTCTGATTTGCTG
 I E E K K V C Q G T S N K L T Q L G T F E D H F L S I Q R N Y D L S F L K T I Q E V A G Y V L I A L I




Figure. Gene Sequencing (Extract)




Figure. SDS-PAGE

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.