

EPC744Ra61 50µg

Eukaryotic Peptidylglycine Alpha Amidating Monooxygenase (PAM)

Organism Species: Rattus norvegicus (Rat)

Instruction manual

FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)



[PROPERTIES]

Source: Eukaryotic expression

Host: 293F Cell

Residues: Phe36~Val820

Tags: N-terminal His Tag

Subcellular Location: Cytoplasm

Purity: > 95%

Traits: Freeze-dried powder

Buffer formulation: PBS, pH7.4, containing 5% Trehalose.

Original Concentration: 400µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 5.7

Predicted Molecular Mass: 77.7kDa

Accurate Molecular Mass: 88kDa as determined by SDS-PAGE reducing conditions.

Phenomenon explanation:

The possible reasons that the actual band size differs from the predicted are as follows:

- 1. Splice variants: Alternative splicing may create different sized proteins from the same gene.
- 2. Relative charge: The composition of amino acids may affects the charge of the protein.
- 3. Post-translational modification: Phosphorylation, glycosylation, methylation etc.
- 4. Post-translation cleavage: Many proteins are synthesized as pro-proteins, and then cleaved to give the active form.
- 5. Polymerization of the target protein: Dimerization, multimerization etc.

[USAGE]

Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.



Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCE]

FKETTRSFSNECLGTIGPVTPLDASDFALDIRMPGVTPKESDTYFCMSMRLPVDEEAFVIDFKPRASMD
TVHHMLLFGCNMPSSTGSYWFCDEGTCTDKANILYAWARNAPPTRLPKGVGFRVGGETGSKYFVLQVHY
GDISAFRDNHKDCSGVSVHLTRVPQPLIAGMYLMMSVDTVIPPGEKVVNADISCQYKMYPMHVFAYRVH
THHLGKVVSGYRVRNGQWTLIGRQNPQLPQAFYPVEHPVDVTFGDILAARCVFTGEGRTEATHIGGTSS
DEMCNLYIMYYMEAKYALSFMTCTKNVAPDMFRTIPAEANIPIPVKPDMVMMHGHHKEAENKEKSALMQ
QPKQGEEEVLEQGDFYSLLSKLLGEREDVHVHKYNPTEKTESGSDLVAEIANVVQKKDLGRSDAREGAE
HEEWGNAILVRDRIHRFHQLESTLRPAESRAFSFQQPGEGPWEPEPSGDFHVEEELDWPGVYLLPGQVS
GVALDSKNNLVIFHRGDHVWDGNSFDSKFVYQQRGLGPIEEDTILVIDPNNAEILQSSGKNLFYLPHGL
SIDTDGNYWVTDVALHQVFKLDPHSKEGPLLILGRSMQPGSDQNHFCQPTDVAVEPSTGAVFVSDGYCN
SRIVQFSPSGKFVTQWGEESSGSSPRPGQFSVPHSLALVPHLDQLCVADRENGRIQCFKTDTKEFVREI
KHASFGRNVFAISYIPGFLFAVNGKPYFGDQEPVQGFVMNFSSGEIIDVFKPVRKHFDMPHDIVASEDG
TVYIGDAHTNTVWKFTLTEKMEHRSV

[IDENTIFICATION]




Figure. Gene Sequencing (Extract)






Figure. SDS-PAGE

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.