

**PAA068Hu01**

**Polyclonal Antibody to Glial Fibrillary Acidic Protein (GFAP)**

**Organism Species: *Homo sapiens* (Human)**

***Instruction manual***

FOR RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

---

13th Edition (Revised in Aug, 2023)

**[ PROPERTIES ]**

**Source:** Polyclonal antibody preparation

**Host:** Rabbit

**Purification:** Antigen-specific affinity chromatography followed by Protein A affinity chromatography

**Traits:** Liquid

**Concentration:** 0.32mg/mL

**UOM:** 100µL

**Cross Reactivity:** Mouse;Rat;Porcine

**Applications:** WB; IHC; FCM

**[ IMMUNOGEN ]**

**Immunogen:** Recombinant GFAP (Met1~Met432) expressed in *E.coli*

**Accession No.:** RPA068Hu01

**[ APPLICATIONS ]**

Western blotting: 0.01-2µg/mL;

Immunohistochemistry: 5-20µg/mL;

Flow cytometry: 20µg/mL

Optimal working dilutions must be determined by end user.

**[ FORMULATION ]**

**Form & Buffer:** Supplied as solution form in 0.01M PBS, pH7.4, containing 0.05% Proclin-300, 50% glycerol.

**[ STORAGE AND STABILITY ]**

**Storage:** Avoid repeated freeze/thaw cycles.


Store at 4°C for frequent use.

Aliquot and store at -20°C for 24 months.


**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no

obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.


**[ IDENTIFICATION ]**




Western Blot; Sample: Porcine Liver lysate Primary Ab: 0.5µg/ml Rabbit Anti-Human GFAP Antibody Second Ab: 0.2µg/ml HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody (Catalog: SAA544Rb19)




DAB staining on IHC-P; Sample: Rat Cerebrum Tissue Primary Ab: 10µg/ml Rabbit Anti-Human GFAP Antibody Control: Used PBS instead of primary antibody Second Ab: 2?g/ml HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody (Catalog: SAA544Rb19)




DAB staining on IHC-P; Sample: Rat Cerebellum Tissue Primary Ab: 10µg/ml Rabbit Anti-Human GFAP Antibody Control: Used PBS instead of primary antibody Second Ab: 2?g/ml HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody (Catalog: SAA544Rb19)



DAB staining on IHC-P; Sample: Rat Spinal cord Tissue Primary Ab: 10µg/ml Rabbit Anti-Human GFAP Antibody Control: Used PBS instead of primary antibody Second Ab: 2?g/ml HRP-Linked



Western Blot; Samples: Lane1: Mouse Cerebrum lysate; Lane2: Rat Cerebrum lysate; Lane3: U87MG cell lysate; Lane4: Porcine Cerebrum lysate; Primary Ab: 0.05µg/ml Rabbit Anti-Human GFAP Antibody Second Ab: 0.2?g/ml HRP-Linked




Western Blot; Samples: Lane1: Rat Cerebrum lysate; Lane2: U87MG cell lysate; Lane3: HBMEC cell lysate; Primary Ab: 0.05µg/ml Rabbit Anti-Human GFAP Antibody Second Ab: 0.2?g/ml HRP-Linked Caprine Anti-Rabbit IgG Polyclonal

Caprine Anti-Rabbit IgG Polyclonal  
Antibody  
(Catalog: SAA544Rb19)

Caprine Anti-Rabbit IgG Polyclonal  
Antibody  
(Catalog: SAA544Rb19)

Antibody  
(Catalog: SAA544Rb19)



Human U87MG cell was fixed with 2% paraformaldehyde (10 min) , permeabilised with 0.1% BSA-Triton X-100, then stained with 20µg/ml rabbit Anti-human GFAP Polyclonal Antibody (Catalog PAA068Hu01, red histogram) or Isotype control antibody (Catalog IS067-Rb01, green histogram), followed by 1µg/ml FITC-conjugated Anti-rabbit IgG Secondary Antibody (Catalog SAA544Rb18).

**[ IMPORTANT NOTE ]**

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.