

**PAB776Hu01**

**Polyclonal Antibody to Interferon Regulatory Factor 8 (IRF8)**

**Organism Species: *Homo sapiens (Human)***

***Instruction manual***

FOR RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

---

13th Edition (Revised in Aug, 2023)

**[ PROPERTIES ]**

**Source:** Polyclonal antibody preparation

**Host:** Rabbit

**Purification:** Antigen-specific affinity chromatography followed by Protein A affinity chromatography

**Traits:** Liquid

**Concentration:** 0.44mg/mL

**UOM:** 100µL

**Cross Reactivity:** N/A

**Applications:** WB; FCM

**[ IMMUNOGEN ]**

**Immunogen:** Recombinant IRF8 (Gly128~Val426) expressed in *E.coli*

**Accession No.:** RPB776Hu01

**[ APPLICATIONS ]**

Western blotting: 0.01-2µg/mL;

Flow cytometry: 20µg/mL;

Optimal working dilutions must be determined by end user.

**[ FORMULATION ]**

**Form & Buffer:** Supplied as solution form in PBS, pH7.4, containing 0.02% NaN<sub>3</sub>, 50% glycerol.

**[ STORAGE AND STABILITY ]**

**Storage:** Avoid repeated freeze/thaw cycles.



Store at 4°C for frequent use.

Aliquot and store at -20°C for 24 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the

expiration date under appropriate storage condition.

**[ IDENTIFICATION ]**



Western Blot; Sample: Human Raji cell Raji human Burkitt's lymphoma cell line lysate; was stained with rabbit Anti-human Primary Ab: 2µg/ml Rabbit Anti-Human IRF8 Polyclonal Antibody (Catalog # PAB776Hu01, filled red histogram) or Second Ab: 0.2µg/mL HRP-Linked Isotype control antibody (Catalog # Caprine Anti-Rabbit IgG Polyclonal Antibody IS067-Rb01, filled green histogram), (Catalog: SAA544Rb19) followed by APC-conjugated Anti-rabbit IgG Secondary Antibody (Catalog # SAA544Rb15). Cells were fixed with 4% paraformaldehyde and permeabilized with 0.1% Triton X-100.

**[ IMPORTANT NOTE ]**

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.