



CSI904Rb01

Primary Rabbit Trigeminal Satellite Glial Cells (SGC)

Organism Species: *Oryctolagus cuniculus* (Rabbit)

Instruction manual

FOR RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

1st Edition (Revised in Aug, 2026)

[DESCRIPTION]

Cell Type: Satellite Glial Cell

Synonyms: SGC

Strain: New Zealand white rabbit

Age: 1-2 Days

Tissue Source: Trigeminal ganglion

Disease: Normal

Size: $>5 \times 10^5$ cells/vial

Growth properties: Adherent

Morphology: Polygonal

[PROPERTIES]

Cell activity: $>85\%$ (Viability by Trypan Blue Exclusion).

Formulation: Frozen 1 mL or T25 flask.

Biosafety: Negative for HIV-1, HBV, HCV, mycoplasma, bacteria, yeast and fungi.

Applications: For research use only. It is not approved for human or animal use, or for application in clinical diagnostic procedures.

[CONTENTS]

Form & Buffer: Supplied as solution form in frozen stock solution (Protein-free, Chemical Defined Cell Cryopreservation Medium). **SGC are cryopreserved at P1 and delivered frozen.**

[USAGE]

Upon receiving the cells in a T-25 flask at room temperature, immediately transfer the cells to 37°C, 5% CO₂ incubator; the cells in vials, directly and immediately transfer the cells from dry ice to liquid nitrogen.

Culture conditions:

DMEM/F12+10% FBS+1% Satellite Glial Cells Growth Supplement+1% Penicillin-Streptomycin Solution

Temperature: 37°C

Condition: 95% air, 5% carbon dioxide

Medium Renewal: Every 1 to 2 days

Dissociation Solution: 0.25% Trypsin

Cell recovery:

After receiving the cells, shake at 37°C in a water bath until completely dissolved, transfer to a 15 ml centrifuge tube, add 3-5 times complete culture solution, 1000 rpm for 5 min, discard the supernatant, and place in a T25 flask for culture.

Cell passage:

1. Cell passage when cell growth at 85-95%.
2. Discard the medium and wash with PBS 1-2 times.
3. Add 1 ml of Trypsin at 37°C, observe the cell under the microscope. If the cells are retracted and rounded, pat the culture flask to let the cells fall off. Stop digestion by adding 2 ml of complete medium containing 10% serum. Make it a single cell suspension.
4. Add the fresh medium to resuspend the cells. Unless otherwise stated, the recommended ratio of primary cells is 1/2-1/3.

[Shipping]

Dry ice.

[STORAGE]

Upon receiving, directly and immediately transfer the cells from dry ice to liquid nitrogen and keep the cells in liquid nitrogen until they are needed for experiments.

[IMPORTANT NOTE]

1. The culture cycle of Primary Rabbit Trigeminal Satellite Glial Cells (SGC) is limited in vitro. It is recommended to use the specialized growth medium provided by Cloud-Clone Corp. and follow the correct operational procedures to ensure optimal culture conditions for these cells.
2. The cell is for research use only, and we will not be responsible for any issue if the cell was used in clinical diagnostic or any other procedures.


[Figure]

Figure 1




Figure 2

Figure 1 Morphology of Rabbit Trigeminal Ganglion Astrocytes (Optical microscope, ×100)

Figure 2 Morphology of Rabbit Trigeminal Ganglion Astrocytes (Optical microscope,×200)




Figure 3

Figure 3 Immunofluorescence identification of GFAP specific antibody (×200)




Figure 4

Figure 4 Immunofluorescence identification of GFAP specific antibody (×400)