

FOR RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

## Basic information

**Product No. :** TSI501Ra49**Organism Species :** Articular Cartilages (Rat) *Same name, Different species.***Strain :** Rat**Sex :** Male**Age :** 8-10 weeks**Weight :** 220g~250g**Level :** SPF**Disease :** Normal Adult**Source :** Articular Cartilage**Traits :** Paraffin block**Applications :** Paraffin slides for pathologic research: IHC,IF and HE,Masson and other stainings.

## BIOSAFETY of the Rat Articular Cartilage Tissue of Normal Adult

Tissues are negative for HIV-1, HBV, HCV, mycoplasma, bacteria, yeast and fungi.

## USAGE of the Rat Articular Cartilage Tissue of Normal Adult

Upon receiving the fresh tissues, transfer the tissues to liquid nitrogen, -80° C freezer or directly process the tissues and experiment immediately. Paraffin embedding tissue often needs to be cut into slices before use. Upon receiving the paraffin section, after grilled with 60° C of 3-5 hours, it can be stored at 4° C in the short term, otherwise it must be stored at -20° C. Upon receiving the frozen section, immediately transfer it to -80° C or -20° C freezer.

## STORAGE of the Rat Articular Cartilage Tissue of Normal Adult

Fresh tissues to be used within 1 month must be stored at -80° C. Paraffin embedding tissue and paraffin section to be used within 1 month may be stored at room temperature, otherwise it must be stored at -4° C(≤3 month) or -20° C(≤6 months). Frozen section must be stored at -20° C(≤3 month) or -80° C(≤6 months). Avoid repeated freeze-thaw cycles.

## FIGURE of the Rat Articular Cartilage Tissue of Normal Adult




Figure 1




Figure 2

Figure 1 Detection of Rat Articular Cartilage Tissue of Normal Adult by HE Staining (Optical microscope,×100)

Figure 2 Detection of Rat Articular Cartilage Tissue of Normal Adult by HE Staining (Optical microscope,×200)