

**RPG487Hu01 1mg**

**Recombinant Farnesyltransferase Alpha (FNTa)**

**Organism Species: *Homo sapiens (Human)***

***Instruction manual***

FOR RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

---

13th Edition (Revised in Aug, 2023)

**[ PROPERTIES ]**

**Source:** Prokaryotic expression

**Host:** *E.coli*

**Residues:** Met51~Lys366

**Tags:** N-terminal His Tag

**Subcellular Location:** Cytoplasm

**Purity:** > 90%

**Traits:** Freeze-dried powder

**Buffer formulation:** PBS, pH7.4, containing 0.01% Sarcosyl, 5% Trehalose.

**Original Concentration:** 1200µg/mL

**Applications:** Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

**Predicted isoelectric point:** 5.1

**Predicted Molecular Mass:** 41.7KDa

**Accurate Molecular Mass:** 41KDa as determined by SDS-PAGE reducing conditions.

**[ USAGE ]**

Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

**[ STORAGE AND STABILITY ]**

**Storage:** Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

**[ SEQUENCE ]**

MDDGFVSLDS PSYVLYRDRA EWADIDPVPQ NDGPNPWWQI IYSDKFRDVI  
DYFRAVLQRD ERSERAFKLT RDAIELNAAN YTVWHFRRVL LKSLQKDLHE  
EMNYITAIIE EQPKNYQVWH HRRVLVEWLR DPSQELEFIA DILNQDAKNI  
HAWQHRQWVI QEFKLWDNEL QYVDQLLKED VRNNSVWNQR YFVISNTTGY  
NDRAVLEREV QYTLEMICLV PHNESAWNYL KGILQDRGLS KYPNLLNQLL  
DLQPSHSSPY LIAFLVDIYE DMLNQCDNK EDILNKALEL CEILAKEKDT  
IRKEYWRYIG RSLQSK

**[ IDENTIFICATION ]**




Figure. SDS-PAGE

**[ IMPORTANT NOTE ]**

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.