

RPG512Hu01 200µg

Recombinant Steroidogenic Acute Regulatory Protein (STAR)

Organism Species: Homo sapiens (Human)

Instruction manual

FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)



[PROPERTIES]

Source: Prokaryotic expression

Host: E.coli

Residues: Met1~Cys285

Tags: N-terminal His and GST Tag

Subcellular Location: Mitochondrion

Purity: > 80%

Traits: Freeze-dried powder

Buffer formulation: PBS, pH7.4, containing 0.01% Sarcosyl, 5% Trehalose.

Original Concentration: 50µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 9.2

Predicted Molecular Mass: 61.9kDa

Accurate Molecular Mass: 62kDa as determined by SDS-PAGE reducing conditions.

[USAGE]

Reconstitute in ddH₂O to a concentration of 0.1-0.2 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCE]



| MLLATFKLCA | GSSYRHMRNM | KGLRQQAVMA | ISQELNRRAL | GGPTPSTWIN |
|------------|------------|------------|------------|------------|
| QVRRRSSLLG | SRLEETLYSD | QELAYLQQGE | EAMQKALGIL | SNQEGWKKES |
| QQDNGDKVMS | KVVPDVGKVF | RLEVVVDQPM | ERLYEELVER | MEAMGEWNPN |
| VKEIKVLQKI | GKDTFITHEL | AAEAAGNLVG | PRDFVSVRCA | KRRGSTCVLA |
| GMATDFGNMP | EQKGVIRAEH | GPTCMVLHPL | AGSPSKTKLT | WLLSIDLKGW |
| LPKSTINOVI | SOTOVDEANH | LRKRLESHPA | SEARC | |

[IDENTIFICATION]




Figure. SDS-PAGE

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.